

# *nodules* on LIDC-IDRI

## I. OVERALL STATISTICS

The confusion matrix for each type is shown in Table I. The overall classification rate is 88.2%. Results for all cases are shown. Scores are presented on the most right of the image. It should be noted that cases not correctly classified are labeled out using red rectangles.

TABLE I  
CONFUSION MATRIX FOR *nodules* ON LIDC-IDRI


	<b>G</b>	<b>W</b>	<b>N</b>	<b>P</b>	<b>V</b>	<b>J</b>
<b>G</b>	<b>0.80</b>	0.01	0.20	0.00	0.00	0.00
<b>W</b>	0.01	<b>0.97</b>	0.03	0.00	0.00	0.0
<b>N</b>	0.00	0.01	<b>0.98</b>	0.01	0.00	0.01
<b>P</b>	0.01	0.00	0.14	<b>0.86</b>	0.00	0.00
<b>V</b>	0.01	0.01	0.05	0.00	<b>0.94</b>	0.00
<b>J</b>	0.00	0.00	0.21	0.00	0.01	<b>0.79</b>



G = Ground glass optic    W = Well-circumscribed    N = Non-nodule  
P = Pleural-tail            V = Vascularized            J = Juxta-pleural


## II. RESULTS

Click highlighted link to jump to corresponding section.



- **II-A** Well-circumscribed
- **II-B** GGO
- **II-C** Juxta-pleural
- **II-D** Pleural-tail
- **II-E** Vascularized
- **II-F** Non-nodule



A. *Well-circumscribed for nodulecircles*









## B. GGO for nodulecircles








## C. Juxta-pleural for nodules/circles
















## D. Pleural-tail for nodulecircles










## E. Vascularized for nodulecircles







F. Non-nodule for nodulecircles

